

MA 1505 Mathematics I  
Tutorial 10 Solutions

1. We have

$$\begin{aligned} f(\mathbf{r}(u, v)) &= f(2u + v, u - 2v, u + 3v) = 2u + v + u - 2v + u + 3v = 4u + 2v, \\ \mathbf{F}(\mathbf{r}(u, v)) &= \mathbf{F}(2u + v, u - 2v, u + 3v) = (2u + v)^2 \mathbf{i} + (u - 2v)^2 \mathbf{j} + (u + 3v)^2 \mathbf{k}, \\ \mathbf{r}_u \times \mathbf{r}_v &= (2\mathbf{i} + \mathbf{j} + \mathbf{k}) \times (\mathbf{i} - 2\mathbf{j} + 3\mathbf{k}) = 5\mathbf{i} - 5\mathbf{j} - 5\mathbf{k}, \\ ||\mathbf{r}_u \times \mathbf{r}_v|| &= 5\sqrt{3}. \end{aligned}$$

Thus if  $R$  denotes the rectangular region

$$0 \leq u \leq 1, \quad 0 \leq v \leq 2,$$

$$\begin{aligned} \int \int_S f(x, y, z) dS &= \int \int_R f(\mathbf{r}(u, v)) ||\mathbf{r}_u \times \mathbf{r}_v|| dA \\ &= \int_0^2 \int_0^1 (4u + 2v)(5\sqrt{3}) dudv = 40\sqrt{3}, \end{aligned}$$

and

$$\begin{aligned} \int \int_S \mathbf{F}(x, y, z) \bullet d\mathbf{S} &= \int \int_R \mathbf{F}(\mathbf{r}(u, v)) \bullet (\mathbf{r}_u \times \mathbf{r}_v) dA \\ &= \int \int_R ((2u + v)^2 \mathbf{i} + (u - 2v)^2 \mathbf{j} + (u + 3v)^2 \mathbf{k}) \bullet (5\mathbf{i} - 5\mathbf{j} - 5\mathbf{k}) dA \\ &= \int_0^2 \int_0^1 (5(2u + v)^2 - 5(u - 2v)^2 - 5(u + 3v)^2) dudv \\ &= \int_0^2 \int_0^1 10(u^2 + uv - 6v^2) dudv = \int_0^2 -\frac{5}{3}(36v^2 - 3v - 2) dv = -\frac{430}{3} \end{aligned}$$

2.  $z = 4 - x^2 - y^2$  parametrizes as  $r(u, v) = u\mathbf{i} + v\mathbf{j} + (4 - u^2 - v^2)\mathbf{k}$ .

$$\mathbf{r}_u \times \mathbf{r}_v = (\mathbf{i} - 2u\mathbf{k}) \times (\mathbf{j} - 2v\mathbf{k}) = 2u\mathbf{i} + 2v\mathbf{j} + \mathbf{k} \implies ||\mathbf{r}_u \times \mathbf{r}_v|| = \sqrt{4u^2 + 4v^2 + 1}$$

$$\text{So } \int \int_S z dS = \int \int_S (4 - u^2 - v^2) dS = \int \int_R (4 - u^2 - v^2) \sqrt{4u^2 + 4v^2 + 1} dA.$$

$R$  is the projection of  $S$  on the  $xy$  plane, which is the circular disk of radius 2.

Use polar coordinates:

$$R : \quad 0 \leq r \leq 2, \quad 0 \leq \theta \leq 2\pi,$$

$$\int \int_R (4 - x^2 - y^2) \sqrt{4x^2 + 4y^2 + 1} dA = \int_0^{2\pi} \int_0^2 (4 - r^2) \sqrt{4r^2 + 1} r dr d\theta \quad (**)$$

$$= 2\pi \left[ \frac{1}{120} (4r^2 + 1)^{3/2} (41 - 6r^2) \right]_0^2 = \frac{289}{60} \pi \sqrt{17} - \frac{41}{60} \pi.$$

(\*\*) To integrate

$$\int (4 - r^2) \sqrt{4r^2 + 1} r dr,$$

you may use the substitution  $u = 4r^2 + 1$ .

3.

$$3x + 2y + z = 6 \implies z = 6 - 3x - 2y,$$

So the surface  $S$  can be represented parametrically as

$$\mathbf{r}(u, v) = u\mathbf{i} + v\mathbf{j} + (6 - 3u - 2v)\mathbf{k}.$$

$$\mathbf{r}_u \times \mathbf{r}_v = (\mathbf{i} - 3\mathbf{k}) \times (\mathbf{j} - 2\mathbf{k}) = 3\mathbf{i} + 2\mathbf{j} + \mathbf{k}$$

The projection of  $S$  on the  $xy$  plane is the region  $R$  bounded by the  $x$ -axis,  $y$ -axis and the line  $3x + 2y = 6$  or  $y = (6 - 3x)/2$ :

$$R : 0 \leq v \leq \frac{6 - 3u}{2}, \quad 0 \leq u \leq 2.$$

The required integral is

$$\begin{aligned} \iint_S \mathbf{F} \bullet d\mathbf{S} &= \iint_R \mathbf{F}(\mathbf{r}(u, v)) \bullet (\mathbf{r}_u \times \mathbf{r}_v) \, dA \\ &= \iint_R (v\mathbf{i} + u^2\mathbf{j} + (6 - 3u - 2v)^2\mathbf{k}) \bullet (3\mathbf{i} + 2\mathbf{j} + \mathbf{k}) \, dA = \int_0^2 \int_0^{(6-3u)/2} (3v + 2u^2 + (6 - 3u - 2v)^2) \, dv \, du \\ &= \int_0^2 \frac{1}{8} (-60u^3 + 291u^2 - 540u + 396) \, du = 31 \end{aligned}$$

4. Let  $\mathbf{F}(x, y, z) = \frac{1}{2}y^2\mathbf{i} + z\mathbf{j} + x\mathbf{k}$ .

By Stoke's Theorem, the given integral is equal to  $\iint_S \operatorname{curl} \mathbf{F} \bullet d\mathbf{S}$ , where  $S$  is the surface on the plane  $x + z = 0$  with  $C$  as boundary.

Substitute  $z = -x$  into  $x^2 + 2y^2 + z^2 = 1$  to get

$$x^2 + 2y^2 + (-x)^2 = 1 \iff x^2 + y^2 = \frac{1}{2}.$$

The projection of  $C$  onto the  $xy$  plane is a circle centred at origin with radius  $1/\sqrt{2}$ .

$$\operatorname{curl} \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{1}{2}y^2 & z & x \end{vmatrix} = -\mathbf{i} - \mathbf{j} - y\mathbf{k}.$$

$S$  can be described as

$$\mathbf{r}(u, v) = u\mathbf{i} + v\mathbf{j} + (-u)\mathbf{k}$$

with

$$\mathbf{r}_u \times \mathbf{r}_v = (\mathbf{i} - \mathbf{k}) \times \mathbf{j} = \mathbf{i} + \mathbf{k}.$$

and  $R$  is the plane region enclosed by  $C$ :

$$R : 0 \leq r \leq \frac{1}{\sqrt{2}}, \quad 0 \leq \theta \leq 2\pi.$$

Hence the required answer is

$$\begin{aligned}
 \iint_S \operatorname{curl} \mathbf{F} \bullet d\mathbf{S} &= \iint_R (-\mathbf{i} - \mathbf{j} - v\mathbf{k}) \bullet (\mathbf{i} + \mathbf{k}) \, dA \\
 &= - \iint_R (1 + v) \, dA = - \int_0^{2\pi} \int_0^{\frac{1}{\sqrt{2}}} (1 + r \sin \theta) \, r \, dr \, d\theta \\
 &= - \int_0^{2\pi} \left[ \frac{1}{4} + \frac{1}{6\sqrt{2}} \sin \theta \right] \, d\theta \\
 &= - \left[ \frac{1}{4}\theta - \frac{1}{6\sqrt{2}} \cos \theta \right]_0^{2\pi} = -\frac{\pi}{2}.
 \end{aligned}$$

5. Using Stoke's Theorem:

Let  $C$  be given by

$$\mathbf{r}(\theta) = \frac{1}{2} \cos \theta \mathbf{i} + \frac{1}{2} \sin \theta \mathbf{j} + \frac{1}{2} \mathbf{k}, \quad (0 \leq \theta \leq 2\pi).$$

Then

$$\mathbf{r}'(\theta) = -\frac{1}{2} \sin \theta \mathbf{i} + \frac{1}{2} \cos \theta \mathbf{j}.$$

Note that the orientation of  $C$  given by this vector equation is anticlockwise. In order to match the orientation of  $S$  given by the outer normal vector, we need to take the negative orientation. Therefore, we will integrate from  $2\pi$  to 0 in the line integral.

$$\mathbf{F}(\mathbf{r}) = \frac{1}{2} \sin \theta \mathbf{i} - \frac{1}{2} \cos \theta \mathbf{j} + \frac{1}{4} \sin \theta \mathbf{k},$$

so that

$$\begin{aligned}
 \iint_S \operatorname{curl} \mathbf{F} \bullet d\mathbf{S} &= \oint_C \mathbf{F}(\mathbf{r}) \bullet d\mathbf{r} \\
 &= \int_{2\pi}^0 -\frac{1}{4} (\sin^2 \theta + \cos^2 \theta) \, d\theta = \int_{2\pi}^0 -\frac{1}{4} \, d\theta = \frac{\pi}{2}.
 \end{aligned}$$

6. By the divergence theorem,

$$\iint_S \mathbf{F} \bullet d\mathbf{S} = \iiint_D \operatorname{div} \mathbf{F} \, dV$$

where  $D$  is the rectangular region given by

$$D : \quad 0 \leq x \leq 1, \quad 0 \leq y \leq 2, \quad -3 \leq z \leq 0$$

and

$$\operatorname{div} \mathbf{F} = \frac{\partial}{\partial x}(x^2) + \frac{\partial}{\partial y}(xy) + \frac{\partial}{\partial z}(x^3y^3) = 3x.$$

So

$$\begin{aligned}\iint_S \mathbf{F} \bullet d\mathbf{S} &= \int_0^1 \int_0^2 \int_{-3}^0 3x \ dz dy dx \\ &= 3 \int_0^1 x \ dx \int_0^2 dy \int_{-3}^0 dz \\ &= 3 \left(\frac{1}{2}\right)(2)(3) = 9\end{aligned}$$